

Product Sheet



QVQ

QUALITY IN ANTIBODIES

Yalelaan 1
3584 CL Utrecht
The Netherlands
+31 30 253 3421


www.qvquality.com
KvK: 30274082
VAT: 8215.17.168
NL88 RABO0153194936

Epithelial cellular adhesion molecule (EpCAM)

Catalogue no.: Q66c
Clone name: QEP-3C7

Product: VHH directed against EpCAM

Target: Epithelial cellular adhesion molecule (EpCAM, UniProtKB P16422), also known as TROP-1, is a 40 kDa type I transmembrane glycoprotein involved in Ca²⁺-independent cell adhesion, signaling, migration and proliferation. EpCAM was originally identified as a tumor-associated antigen because of its expression in a large number of cancers. EpCAM is expressed in basolateral cell membranes of epithelial cells. The protein consists of an 242 amino acid ectodomain that contains a signal peptide (SP) and two EGF-like subdomains, a 23 amino acid transmembrane domain and a cytoplasmic domain of only 26 amino acids. The 2nd EGF-like domain is also referred to as a thyroglobulin-like domain. This cytoplasmic domain of EpCAM interacts directly with α -actinin of the actin cytoskeleton.¹⁻⁴



Source: Recombinant monoclonal VHH (Llama glama), purified from *S.cerevisiae* using affinity chromatography. Immunization with cancer cells. Phage-display selection on recombinant protein using total elution.

Specificity: Human EpCAM.


Formulation: 0.2 μ m filtered solution in PBS.

Mol. Weight: 15.3 kDa
Ext. Coeff. (ϵ): 24535
A₂₈₀ at 1g/L: 1.6

Storage: Shipped on blue ice. Store at 4 °C or -20 °C (aliquots). Addition of 0.02% sodiumazide is optional.

Applications: ELISA

Examples:



Binding of Q66c to recombinant EpCAM in ELISA.

References:

- 1 Huang et al. (2018) *Int J Mol Med.* 42, 1771-1785
- 2 Balzar et al. (1999) *J Mol Med.* 77, 699-712
- 3 Schnell et al. (2013) *Biochim Biophys Acta.* 1828, 1889-2001
- 4 Balzar et al. (1998) *Mol Cell Biol.* 18, 4833-4843