

Product Sheet



QVQ

QUALITY IN ANTIBODIES

Yalelaan 1
3584 CL Utrecht
The Netherlands
+31 30 253 3421


www.qvquality.com
KvK: 30274082
VAT: 8215.17.168
NL88 RABO0153194936

Urokinase Plasminogen Activator Surface receptor (uPAR)

Catalogue no.: Q88c
Clone name: QUP-2G4

Product: VHH directed against uPAR

Target: Urokinase plasminogen activator surface receptor (uPAR, UniProtKB Q03405), also known as monocyte activation antigen Mo3, PLAUR and CD87, is a GPI-anchored protein that binds the urokinase-type Plasminogen Activator (uPA). uPAR is a 335 amino acid large GPI-anchored protein and functions by localizing and enhancing the uPA-mediated conversion of plasminogen to plasmin. In addition, uPAR can interact with vitronectin and integrins and hereby regulate cell adhesion. Ligand binding to uPAR results in downstream signal transduction towards tyrosine kinases, gene expression, cell adhesion, and chemotaxis. uPAR is expressed a large number of cancers and is linked to tumor metastasis and invasion.¹⁻³



Source: Recombinant monoclonal VHH (Llama glama), purified from *S. cerevisiae* using affinity chromatography. Immunization and phage-display selection on immobilized recombinant human uPAR.

Specificity: Domains D2-D3 of human uPAR.

Formulation: 0.2 µm filtered solution in PBS.

Mol. Weight: 15.0 kDa


Ext. Coeff. (ε): 25565

A₂₈₀ at 1g/L: 1.7

Storage: Shipped on blue ice. Store at 4 °C or -20 °C (aliquots). Addition of 0.02% sodiumazide is optional.

Applications: ELISA

Examples:



Binding of Q88 to recombinant uPAR in ELISA.

References:

- 1 [Ploeg, M. \(2013\) Theranostics, 3:467-476.](#)
- 2 [Dear, AE, and Medcalf, RL. \(1998\) Eur J Biochem, 252:185-193.](#)
- 3 [Boonstra et al. \(2011\) Curr Pharm Des, 17:1890-1910](#)