

Product Sheet



Yalelaan 1
3584 CL Utrecht
The Netherlands
+31 30 253 3421

www.qvquality.com
KvK: 30274082
VAT: 8215.17.168
NL88 RABO0153194936


Programmed Death Ligand 1 (PD-L2)

Catalogue no.: Q90c

Clone name: QL2-1A4

Product: VHH directed against PD-L2

Target: Butyrophilin B7-DC, also called programmed death ligand 2 (PD-L2) (UniProtKB Q9BQ51), is a member of the B7 family of immune proteins that provide signals for both stimulating and inhibiting T cell activation. Other family members include B7-1, B7-2, B7-H2, PD-L1 and B7-H3. B7 proteins are members of the immunoglobulin (Ig) superfamily, their extracellular domains contain 2 Ig-like domains and all members have short cytoplasmic domains. PD-L2 has been identified as one of two ligands for programmed death-1 (PD-1), a member of the CD28 family of immunoreceptors. The PD-L2:PD-1 pathway is involved in the negative regulation of immune responses and in the regulation of peripheral tolerance.¹⁻³



Source: Recombinant monoclonal VHH (Llama glama), purified from *S.cerevisiae* using affinity chromatography. Immunization with cancer cells. Phage-display selection on recombinant PD-L2 using total elution.

Specificity: Human PD-L2.

Formulation: 0.2 µm filtered solution in PBS.

Mol. Weight: 16.0 kDa


Ext. Coeff. (ε): 28545

A₂₈₀ at 1g/L: 1.8

Storage: Shipped on blue ice. Store at 4 °C or -20 °C (aliquots). Addition of 0.02% sodiumazide is optional.

Applications: ELISA

Examples:



Binding of Q90 to recombinant PD-L2 ectodomain in ELISA.

References:

- 1 Greenwald R.J. et al. (2005) *Annu Rev Immunol* 23:515-548
- 2 Lázár-Molnár E. et al. (2008) *Proc Natl Acad Sci U S A*, 105:10483-10488
- 3 Nishimura, H. and T. Honjo (2001) *Trends in Immunology* 22:265